

TABLE S1. Primers for mRNA expression analysis of reproductive and gestational tissue

| Gene | Forward and Reverse Prime Sequence | GeneBank accession # |
|--------------|--|----------------------|
| <i>Il1a</i> | F-5' CCGACCTCATTTCTTCTGG 3' R-5' GTGCACCCGACTTGTTCTT 3' | NM_010554.4 |
| <i>Il1b</i> | F-5' CCAAAGCAATAACCAAAGAAA 3' R-5' GCTTGTGCTCTGCTTGAG 3' | NM 008361.3 |
| <i>Il6</i> | F-5' ACAACCACGGCCTCCCTAC 3' R-5' TCCACGATTTCCCAGAGAAC 3' | NM 031168.1 |
| <i>Il10</i> | F-5' AGGCGCTGTCATCGATTCT 3' R-5' TGGCCTTGTAGACACCTTGGT 3' | NM 010548.2 |
| <i>Tnf</i> | F-5' GTAGCCCACGTCGTA 3' R-5' TCCACGATTTCCCAG 3' | NM 013693.3 |
| <i>Ptgs2</i> | F-5' GTTTGCATTCTTGCCCAGC 3' R-5' AGTCCACTCCATGGCCCAGT 3' | NM 011198.3 |
| <i>Actb</i> | F-5' CGTGGGCCGCCCTAGGCACCA 3' R-5' ACACGCAGCTCATTGTA 3' | NM 007393.3 |

TABLE S2. Effect of maternal LPS and (+)-naloxone treatment on body morphometry (absolute weight) in male and female progeny at 20 weeks.

| Absolute weight | PBS | (+)-naloxone | LPS | LPS / (+)-nal |
|--------------------------|--------------------------|-------------------------|--------------------------|--------------------------|
| <i>Male progeny</i> | | | | |
| N, pups (dams) | 24 (10) | 21 (7) | 14 (5) | 25 (7) |
| Total body weight (g) * | 25.0 ± 0.5 ^a | 27.5 ± 0.6 ^b | 26.6 ± 0.8 ^{ab} | 25.9 ± 0.6 ^a |
| Lean body weight (g) | 24.4 ± 0.4 ^a | 27.1 ± 0.4 ^b | 24.9 ± 0.5 ^a | 25.6 ± 0.4 ^{ab} |
| Total muscle (mg) § | 748 ± 19 ^{ab} | 831 ± 20 ^c | 692 ± 25 ^b | 817 ± 20 ^{ac} |
| Total central fat (mg) ^ | 734 ± 34 ^a | 902 ± 36 ^b | 864 ± 46 ^{ab} | 840 ± 37 ^{ab} |
| Brain (mg) | 413 ± 7 | 433 ± 9 | 422 ± 7 | 411 ± 7 |
| Heart (mg) | 133 ± 4 | 144 ± 4 | 138 ± 5 | 136 ± 4 |
| Lungs (mg) # | 193 ± 6 | 195 ± 6 | 184 ± 8 | 180 ± 6 |
| Kidneys (mg) # | 329 ± 8 | 357 ± 9 | 335 ± 11 | 330 ± 9 |
| Liver (mg) | 1225 ± 33 | 1220 ± 44 | 1335 ± 35 | 1301 ± 36 |
| Adrenals (mg) # | 5.25 ± 0.36 | 5.30 ± 0.49 | 5.10 ± 0.38 | 4.43 ± 0.40 |
| Thymus (mg) | 46.5 ± 2.2 ^a | 57.5 ± 2.4 ^b | 57.9 ± 3.0 ^b | 55.0 ± 2.4 ^{ab} |
| Spleen (mg) | 62.4 ± 4.8 ^a | 83 ± 5 ^b | 72.8 ± 6.5 ^{ab} | 71.6 ± 5.3 ^{ab} |
| Testes (mg) # | 184 ± 5 | 190 ± 5 | 187 ± 7 | 181 ± 5 |
| Seminal vesicle (mg) | 258 ± 11 ^{ab} | 258 ± 12 ^{ab} | 220 ± 16 ^a | 275 ± 13 ^b |
| Epididymis (mg) | 82.4 ± 4.8 ^{ab} | 86 ± 5 ^{ab} | 100 ± 7 ^a | 74.0 ± 5.3 ^b |
| <i>Female progeny</i> | | | | |
| N, pups (dams) | 21 (10) | 8 (5) | 9 (4) | 13 (7) |
| Total body weight (g) * | 21.9 ± 0.5 | 22.8 ± 0.7 | 21.4 ± 0.8 | 21.7 ± 0.6 |
| Lean body weight (g) | 21.0 ± 3.8 | 22.0 ± 6.2 | 20.3 ± 5.8 | 20.7 ± 0.5 |
| Total muscle (mg) § | 587 ± 15 ^{ab} | 661 ± 24 ^a | 541 ± 23 ^b | 608 ± 20 ^{ab} |
| Total central fat (mg) ^ | 716 ± 34 | 618 ± 55 | 650 ± 52 | 628 ± 45 |
| Brain (mg) | 412 ± 6 | 419 ± 9 | 415 ± 9 | 388 ± 8 |
| Heart (mg) | 112 ± 3 | 106 ± 4 | 120 ± 4 | 109 ± 4 |
| Lungs (mg) # | 183 ± 7 | 180 ± 11 | 178 ± 11 | 168 ± 9 |
| Kidneys (mg) # | 262 ± 6 | 276 ± 11 | 254 ± 10 | 256 ± 9 |
| Liver (mg) | 1111 ± 36 | 1020 ± 55 | 1219 ± 59 | 1171 ± 48 |
| Adrenals (mg) # | 6.08 ± 0.52 | 8.21 ± 0.80 | 6.70 ± 0.85 | 5.81 ± 0.69 |
| Thymus (mg) | 59.8 ± 2.9 | 66.6 ± 4.6 | 57.5 ± 4.4 | 65.5 ± 3.8 |
| Spleen (mg) | 69.2 ± 2.7 | 71.0 ± 4.4 | 65.5 ± 4.1 | 70.8 ± 3.6 |
| Uterus (mg) | 59.2 ± 5.2 | 68.2 ± 8.4 | 58.4 ± 12.2 | 59.4 ± 7.2 |
| Ovaries (mg) # | 7.13 ± 0.63 | 7.50 ± 1.03 | 8.46 ± 1.50 | 7.14 ± 0.88 |

* Data are estimated marginal mean ± SEM absolute weights in 20 week old male and female progeny of dams administered LPS and/or (+)-naloxone in late gestation. Effect of treatment group was analysed by Mixed Model Linear Repeated Measures ANOVA, using litter size as covariate.

^ Total central fat is the sum of all fat depots measured. The number of progeny in each group is indicated, with the number of pregnant dams in parentheses. Data on individual fat deposits and muscle tissues are shown in Table 2.

§ Total muscle is the sum of all muscles measured.

Combined weight of left and right organs.

a,b,c Different superscript letters indicate differences between treatment groups ($p < 0.05$).

TABLE S3. Effect of maternal LPS and (+)-naloxone treatment on body composition (relative) in male and female progeny at 20 weeks.

| Relative weight | PBS | (+)-naloxone | LPS | LPS / (+)-nal |
|--------------------------------|---------------------------|---------------------------|---------------------------|---------------------------|
| <i>Male progeny</i> | | | | |
| N, pups (dams) | 24 (10) | 21 (7) | 14 (5) | 25 (7) |
| Total muscle (%) ^{§*} | 3.59 ± 0.09 ^{ab} | 3.99 ± 0.01 ^c | 3.32 ± .012 ^b | 3.93 ± 0.10 ^{ac} |
| Total central fat (%)* | 3.52 ± 0.17 ^a | 4.28 ± 0.22 ^b | 4.32 ± 0.18 ^b | 3.98 ± 0.16 ^{ab} |
| Brain (%) | 1.70 ± 0.03 ^a | 1.56 ± 0.03 ^b | 1.75 ± 0.04 ^a | 1.61 ± 0.03 ^{ab} |
| Heart (%) | 0.64 ± 0.02 | 0.69 ± 0.02 | 0.66 ± 0.02 | 0.65 ± 0.02 |
| Lungs (%) [#] | 0.93 ± 0.03 | 0.94 ± 0.03 | 0.88 ± 0.04 | 0.87 ± 0.03 |
| Kidneys (%) [#] | 1.58 ± 0.04 | 1.73 ± 0.04 | 1.61 ± 0.05 | 1.59 ± 0.04 |
| Liver (%) | 5.88 ± 0.16 | 6.41 ± 0.17 | 5.86 ± 0.21 | 6.25 ± 0.17 |
| Adrenals (%) [#] | 0.03 ± 0.00 | 0.02 ± 0.00 | .03 ± 0.00 | 0.02 ± 0.00 |
| Thymus (%) | 0.22 ± 0.01 ^a | 0.28 ± 0.01 ^b | 0.28 ± 0.01 ^b | 0.27 ± 0.01 ^{ab} |
| Spleen (%) | 0.30 ± 0.02 ^a | 0.40 ± 0.02 ^b | 0.35 ± 0.03 ^{ab} | 0.34 ± 0.03 ^{ab} |
| Testes (%) [#] | 0.88 ± 0.02 | 0.91 ± 0.03 | 0.90 ± 0.03 | 0.87 ± 0.03 |
| Seminal vesicle (%) | 1.24 ± 0.06 ^{ab} | 1.24 ± 0.06 ^{ab} | 1.06 ± 0.08 ^a | 1.32 ± 0.06 ^b |
| Epididymis (%) | 0.40 ± 0.02 ^{ab} | 0.41 ± 0.02 ^{ab} | 0.48 ± 0.03 ^a | 0.36 ± 0.03 ^b |
| <i>Female progeny</i> | | | | |
| N, pups (dams) | 21 (10) | 8 (5) | 9 (4) | 13 (7) |
| Total muscle (%) ^{§*} | 2.82 ± 0.07 ^{ab} | 3.18 ± 0.12 ^a | 2.60 ± 0.11 ^b | 2.92 ± 0.10 ^{ab} |
| Total central fat (%)* | 3.43 ± 0.16 | 2.96 ± 0.26 | 3.11 ± 0.25 | 2.95 ± 0.21 |
| Brain (%) | 1.97 ± 0.04 | 1.91 ± 0.06 | 2.05 ± 0.05 | 1.89 ± 0.05 |
| Heart (%) | 0.54 ± 0.01 | 0.58 ± 0.02 | 0.51 ± 0.02 | 0.52 ± 0.02 |
| Lungs (%) [#] | 0.88 ± 0.03 | 0.87 ± 0.05 | 0.85 ± 0.05 | 0.81 ± 0.04 |
| Kidneys (%) [#] | 1.26 ± 0.03 | 1.32 ± 0.05 | 1.22 ± 0.05 | 1.29 ± 0.04 |
| Liver (%) | 5.34 ± 0.17 | 5.85 ± 0.28 | 4.90 ± 0.26 | 5.62 ± 0.23 |
| Adrenals (%) [#] | 0.03 ± 0.00 | 0.03 ± 0.00 | 0.04 ± 0.00 | 0.03 ± 0.00 |
| Thymus (%) | 0.29 ± 0.01 | 0.32 ± 0.02 | 0.28 ± 0.02 | 0.31 ± 0.02 |
| Spleen (%) | 0.33 ± 0.01 | 0.34 ± 0.02 | 0.31 ± 0.02 | 0.34 ± 0.02 |
| Uterus (%) | 0.28 ± 0.02 | 0.33 ± 0.04 | 0.28 ± 0.06 | 0.29 ± 0.03 |
| Ovaries (%) [#] | 0.03 ± 0.00 | 0.04 ± 0.00 | 0.04 ± 0.01 | 0.03 ± 0.00 |

* Data are estimated marginal mean ± SEM relative weights in 20 week old male and female progeny of dams administered LPS and/or (+)-naloxone in late gestation. Effect of treatment group was analysed by Mixed Model Linear Repeated Measures ANOVA, using litter size as covariate. The number of progeny in each group is indicated, with the number of pregnant dams in parentheses. Data on individual fat deposits and muscle tissues are shown in Table 3.

[^] Total central fat is the sum of all fat depots measured. [§] Total muscle is the sum of all muscles measured.

[#] Combined weight of left and right organs.

^{a,b,c} Different superscript letters indicate differences between treatment groups ($p < 0.05$).